



Childhood immunisation schedule

Use the checklist below to tick off your child's immunisations as you go.

Age	Disease protected against	
Birth	<ul style="list-style-type: none">• Hepatitis B (1 injection) (<i>usually offered in hospital</i>)	<input type="checkbox"/>
6–8 weeks	<ul style="list-style-type: none">• Diphtheria, tetanus, pertussis (whooping cough), hepatitis B, poliomyelitis, haemophilus influenzae type b (1 injection)• Pneumococcal (1 injection)• Rotavirus (drops in mouth)	<input type="checkbox"/>
4 months	As above	<input type="checkbox"/>
6 months	As above	<input type="checkbox"/>
12 months	<ul style="list-style-type: none">• Haemophilus influenzae type b, meningococcal C (1 injection)• Measles, mumps, rubella (1 injection)	<input type="checkbox"/>
The following groups need an additional vaccine at 12 months:		
Aboriginal children	<ul style="list-style-type: none">• Hepatitis A (1 injection)	<input type="checkbox"/>
Pre-term or low birth weight	<ul style="list-style-type: none">• Hepatitis B (1 injection)	<input type="checkbox"/>
Medically at risk	<ul style="list-style-type: none">• Pneumococcal (1 injection)	<input type="checkbox"/>
18 months	<ul style="list-style-type: none">• Measles, mumps, rubella, varicella (1 injection) only for those who received a MMR vaccine previously• Diphtheria, tetanus, pertussis (whooping cough)	<input type="checkbox"/>
18 months Aboriginal children only	<ul style="list-style-type: none">• Pneumococcal (if not already given at 12 months)• Hepatitis A	<input type="checkbox"/>
4 years	<ul style="list-style-type: none">• Diphtheria, tetanus, pertussis (whooping cough), poliomyelitis (1 injection)	<input type="checkbox"/>
Medically at risk	<ul style="list-style-type: none">• Pneumococcal (1 injection)	<input type="checkbox"/>

All children aged from 6 months to less than 5 years should also receive the influenza vaccine each year. WA Health also offers a **FREE** vaccination program for year 8 students provided across 3 terms.